

A level Mathematics Practice Paper – Integration – Mark scheme

| Question | Scheme | Marks |
|-------------|---|-------------------|
| 1(a) | $1 = A(3x-1)^2 + Bx(3x-1) + Cx$ | B1 |
| | $x \rightarrow 0$ $(1 = A)$ | M1 |
| | $x \rightarrow \frac{1}{3}$ $1 = \frac{1}{3}C \Rightarrow C = 3$ any two constants correct | A1 |
| | Coefficients of x^2 | |
| | $0 = 9A + 3B \Rightarrow B = -3$ all three constants correct | A1 |
| | | (4) |
| 1(b) | (i) $\int \left(\frac{1}{x} - \frac{3}{3x-1} + \frac{3}{(3x-1)^2} \right) dx$ | |
| | $= \ln x - \frac{3}{3} \ln(3x-1) + \frac{3}{(-1)3} (3x-1)^{-1} \quad (+C)$ | M1 A1ft A1ft |
| | $\left(= \ln x - \ln(3x-1) - \frac{1}{3x-1} \quad (+C) \right)$ | |
| | (ii) $\int_1^2 f(x) dx = \left[\ln x - \ln(3x-1) - \frac{1}{3x-1} \right]_1^2$ | |
| | $= \left(\ln 2 - \ln 5 - \frac{1}{5} \right) - \left(\ln 1 - \ln 2 - \frac{1}{2} \right)$ | M1 |
| | $= \ln \frac{2 \times 2}{5} + \dots$ | M1 |
| | $= \frac{3}{10} + \ln \left(\frac{4}{5} \right)$ | A1 |
| | | (6) |
| | | (10 marks) |

A level Mathematics Practice Paper – Integration – Mark scheme

| Question | Scheme | Marks | |
|----------------------------------|---|--|-----------|
| 2 | $\int x \sin 2x \, dx = -\frac{x \cos 2x}{2} + \int \frac{\cos 2x}{2} \, dx$ | M1 A1 A1 | |
| | $= \dots + \frac{\sin 2x}{4}$ | M1 | |
| | $\left[\dots \right]_0^{\frac{\pi}{2}} = \frac{\pi}{4}$ | M1 A1 | |
| (6 marks) | | | |
| 3(a) | $\int \frac{1}{x^3} \ln x \, dx, \quad \left\{ \begin{array}{l} u = \ln x \Rightarrow \frac{du}{dx} = \frac{1}{x} \\ \frac{dv}{dx} = x^{-3} \Rightarrow v = \frac{x^{-2}}{-2} = \frac{-1}{2x^2} \end{array} \right\}$ | | |
| | $= \frac{-1}{2x^2} \ln x - \int \frac{-1}{2x^2} \cdot \frac{1}{x} \, dx$ | In the form $\frac{\pm \lambda}{x^2} \ln x \pm \int \mu \frac{1}{x^2} \cdot \frac{1}{x}$ | M1 |
| | | $\frac{-1}{2x^2} \ln x$ simplified or un-simplified. | <u>A1</u> |
| | | $-\int \frac{-1}{2x^2} \cdot \frac{1}{x}$ simplified or un-simplified. | <u>A1</u> |
| | $\left\{ = \frac{-1}{2x^2} \ln x + \frac{1}{2} \int \frac{1}{x^3} \, dx \right\}$ | | |
| | $= -\frac{1}{2x^2} \ln x + \frac{1}{2} \left(-\frac{1}{2x^2} \right) \{+c\}$ | $\pm \int \mu \frac{1}{x^2} \cdot \frac{1}{x} \rightarrow \pm \beta x^{-2}.$ | dM1 |
| Correct answer, with/without + c | | A1 | |
| (5) | | | |
| 3(b) | $\left\{ \left[-\frac{1}{2x^2} \ln x - \frac{1}{4x^2} \right]_1^2 \right\} = \left(-\frac{1}{2(2)^2} \ln 2 - \frac{1}{4(2)^2} \right) - \left(-\frac{1}{2(1)^2} \ln 1 - \frac{1}{4(1)^2} \right)$ | | |
| | Applies limits of 2 and 1 to their part (a) answer and subtracts the correct way round. | M1 | |
| | $= \frac{3}{16} - \frac{1}{8} \ln 2 \quad \text{or} \quad \frac{3}{16} - \ln 2^{\frac{1}{8}} \quad \text{or} \quad \frac{1}{16} (3 - 2 \ln 2), \text{ etc, or awrt } 0.1 \text{ or equivalent.}$ | A1 | |
| (2) | | | |
| (7 marks) | | | |

A level Mathematics Practice Paper – Integration – Mark scheme

| Question | Scheme | Marks |
|-------------|--|------------------|
| 4(a) | $\int x \sin 3x \, dx = -\frac{1}{3}x \cos 3x - \int -\frac{1}{3} \cos 3x \{dx\}$ | M1 A1 |
| | $= -\frac{1}{3}x \cos 3x + \frac{1}{9} \sin 3x \{+ c\}$ | A1 |
| | | (3) |
| 4(b) | $\int x^2 \cos 3x \, dx = \frac{1}{3}x^2 \sin 3x - \int \frac{2}{3}x \sin 3x \{dx\}$ | M1 A1 |
| | $= \frac{1}{3}x^2 \sin 3x - \frac{2}{3} \left(-\frac{1}{3}x \cos 3x + \frac{1}{9} \sin 3x \right) \{+ c\}$ | A1 isw |
| | $\left\{ = \frac{1}{3}x^2 \sin 3x + \frac{2}{9}x \cos 3x - \frac{2}{27} \sin 3x \{+ c\} \right\}$ | |
| | <i>Ignore subsequent working</i> | (3) |
| | | (6 marks) |
| 5(a) | $y = 4x - xe^{\frac{1}{2}x}, x \geq 0$ | |
| | $\left\{ y=0 \quad 4x - xe^{\frac{1}{2}x} = 0 \quad x(4 - e^{\frac{1}{2}x}) = 0 \right\}$ | |
| | $e^{\frac{1}{2}x} = 4 \quad x_A = 4 \ln 2$ | M1 A1 |
| | | (2) |
| 5(b) | $\left\{ xe^{\frac{1}{2}x} dx \right\} = 2xe^{\frac{1}{2}x} \quad 2e^{\frac{1}{2}x} \{dx\}$ | M1 A1 |
| | $= 2xe^{\frac{1}{2}x} \quad 4e^{\frac{1}{2}x} \{+ c\}$ | A1 |
| | | (3) |
| 5(c) | $\left\{ 4x dx \right\} = 2x^2$ | B1 |
| | $\left\{ \int_0^{4 \ln 2} (4x - xe^{\frac{1}{2}x}) dx \right\} = \left[2x^2 - \left(2xe^{\frac{1}{2}x} - 4e^{\frac{1}{2}x} \right) \right]_0^{4 \ln 2 \text{ or } \ln 16 \text{ or their limits}}$ | |
| | $= \left(2(4 \ln 2)^2 - 2(4 \ln 2)e^{\frac{1}{2}(4 \ln 2)} + 4e^{\frac{1}{2}(4 \ln 2)} \right) - \left(2(0)^2 - 2(0)e^{\frac{1}{2}(0)} + 4e^{\frac{1}{2}(0)} \right)$ | M1 |
| | $= (32(\ln 2)^2 - 32(\ln 2) + 16) - (4)$ | |
| | $= 32(\ln 2)^2 - 32(\ln 2) + 12$ | A1 |
| | | (3) |
| | | (8 marks) |

A level Mathematics Practice Paper – Integration – Mark scheme

| Question | Scheme | | Marks |
|-------------|--|--|------------------|
| 6(a) | 1.154701 | | B1 cao |
| | | | (1) |
| 6(b) | Area $\approx \frac{1}{2} \times \frac{\pi}{6} \times [1 + 2(1.035276 + \text{their } 1.154701) + 1.414214]$ | | B1 M1 |
| | $= \frac{\pi}{12} \times 6.794168 = 1.778709023\dots = 1.7787$ (4 dp) | 1.7787 or awrt 1.7787 | A1 |
| | | | (3) |
| 6(c) | $V = \pi \int_0^{\frac{\pi}{2}} \left(\sec\left(\frac{x}{2}\right) \right)^2 dx$ | For $\pi \int \left(\sec\left(\frac{x}{2}\right) \right)^2$. Ignore limits and dx. Can be implied. B1 | |
| | $= \{\pi\} \left[2 \tan\left(\frac{x}{2}\right) \right]_0^{\frac{\pi}{2}}$ | $\pm \lambda \tan\left(\frac{x}{2}\right)$ M1 | |
| | | $2 \tan\left(\frac{x}{2}\right)$ or equivalent A1 | |
| | $= 2\pi$ | 2π A1 cao cso | |
| | | | (4) |
| | | | (8 marks) |

A level Mathematics Practice Paper – Integration – Mark scheme

| Question | Scheme | Marks |
|----------|---|-------------------|
| 7(a) | $\frac{5}{(x-1)(3x+2)} = \frac{A}{x-1} + \frac{B}{3x+2}$ | |
| | $5 = A(3x+2) + B(x-1)$ | |
| | $x \rightarrow 1$ $5 = 5A \Rightarrow A = 1$ | M1 A1 |
| | $x \rightarrow -\frac{2}{3}$ $5 = -\frac{5}{3}B \Rightarrow B = -3$ | A1 |
| | | (3) |
| 7(b) | $\int \frac{5}{(x-1)(3x+2)} dx = \int \left(\frac{1}{x-1} - \frac{3}{3x+2} \right) dx$ | |
| | $= \ln(x-1) - \ln(3x+2) + C$ ft constants | M1 A1ft A1ft |
| | | (3) |
| 7(c) | $\int \frac{5}{(x-1)(3x+2)} dx = \int \left(\frac{1}{y} \right) dy$ | M1 |
| | $\ln(x-1) - \ln(3x+2) = \ln y + C$ | M1 A1 |
| | $y = \frac{K(x-1)}{3x+2}$ depends on first two Ms in (c) | M1 dep |
| | Using (2, 8) $8 = \frac{K}{8}$ depends on first two Ms in (c) | M1 dep |
| | $y = \frac{64(x-1)}{3x+2}$ | A1 |
| | | (6) |
| | | (12 marks) |

A level Mathematics Practice Paper – Integration – Mark scheme

| Question | Scheme | Marks |
|-------------|---|---------------------------|
| 8(a) | $\frac{3y-4}{y(3y+2)} \equiv \frac{A}{y} + \frac{B}{(3y+2)} \Rightarrow 3y-4 = A(3y+2) + By$ $y=0 \Rightarrow -4 = 2A \Rightarrow A = -2$ $y = -\frac{2}{3} \Rightarrow -6 = -\frac{2}{3}B \Rightarrow B = 9$ | M1 A1 A1 |
| | $\int \frac{3y-4}{y(3y+2)} dy = \int \frac{-2}{y} + \frac{9}{(3y+2)} dy$ $= -2\ln y + 3\ln(3y+2) \{+c\}$ | M1 A1 ft A1 cao |
| | | (6) |
| 8(b) | $\{x = 4\sin^2 \theta \Rightarrow\} \frac{dx}{d\theta} = 8\sin \theta \cos \theta \text{ or } \frac{dx}{d\theta} = 4\sin 2\theta \text{ or } dx = 8\sin \theta \cos \theta d\theta$ | B1 |
| | $\int \sqrt{\frac{4\sin^2 \theta}{4-4\sin^2 \theta}} \cdot 8\sin \theta \cos \theta \{d\theta\} \text{ or } \int \sqrt{\frac{4\sin^2 \theta}{4-4\sin^2 \theta}} \cdot 4\sin 2\theta \{d\theta\}$ | M1 |
| | $= \int \underline{\underline{\tan \theta}} \cdot 8\sin \theta \cos \theta \{d\theta\} \text{ or } \int \underline{\underline{\tan \theta}} \cdot 4\sin 2\theta \{d\theta\}$ | <u>M1</u> |
| | $= \int 8\sin^2 \theta d\theta$ | A1 |
| | $3 = 4\sin^2 \theta \text{ or } \frac{3}{4} = \sin^2 \theta \text{ or } \sin \theta = \frac{\sqrt{3}}{2} \Rightarrow \theta = \frac{\pi}{3} \quad \{x=0 \rightarrow \theta=0\}$ | B1 |
| | | (5) |
| 8(c) | $= \{8\} \int \left(\frac{1-\cos 2\theta}{2}\right) d\theta \quad \left\{ = \int (4-4\cos 2\theta) d\theta \right\}$ | M1 |
| | $= \{8\} \left(\frac{1}{2}\theta - \frac{1}{4}\sin 2\theta\right) \quad \{= 4\theta - 2\sin 2\theta\}$ | M1 A1 |
| | $\left\{ \int_0^{\frac{\pi}{3}} 8\sin^2 \theta d\theta = 8 \left[\frac{1}{2}\theta - \frac{1}{4}\sin 2\theta \right]_0^{\frac{\pi}{3}} \right\} = 8 \left(\left(\frac{\pi}{6} - \frac{1}{4} \left(\frac{\sqrt{3}}{2} \right) \right) - (0+0) \right)$ | |
| | $= \frac{4}{3}\pi - \sqrt{3} \text{ o.e.}$ | A1 |
| | | (4) |
| | | (15 marks) |

A level Mathematics Practice Paper – Integration – Mark scheme

| Question | Scheme | Marks | | |
|-------------|---|--|----|-----------------|
| 9(a) | <i>Working parametrically:</i> | | | |
| | $x = 1 - \frac{1}{2}t, \quad y = 2^t - 1 \text{ or } y = e^{t \ln 2} - 1$ | | | |
| | $\{x = 0 \Rightarrow\} 0 = 1 - \frac{1}{2}t \Rightarrow t = 2$ | Applies $x = 0$ to obtain a value for t . | M1 | |
| | When $t = 2, \quad y = 2^2 - 1 = 3$ | Correct value for y . | A1 | |
| | | (2) | | |
| 9(b) | $\{y = 0 \Rightarrow\} 0 = 2^t - 1 \Rightarrow t = 0$ | Applies $y = 0$ to obtain a value for t . (Must be seen in part (b)). | M1 | |
| | When $t = 0, \quad x = 1 - \frac{1}{2}(0) = 1$ | $x = 1$ | A1 | |
| | | (2) | | |
| 9(c) | $\frac{dx}{dt} = -\frac{1}{2}$ and either $\frac{dy}{dt} = 2^t \ln 2$ or $\frac{dy}{dt} = e^{t \ln 2} \ln 2$ | | B1 | |
| | $\frac{dy}{dx} = \frac{2^t \ln 2}{-\frac{1}{2}}$ | Attempts their $\frac{dy}{dt}$ divided by their $\frac{dx}{dt}$. | M1 | |
| | At A, $t = "2"$, so $m(\mathbf{T}) = -8 \ln 2 \Rightarrow m(\mathbf{N}) = \frac{1}{8 \ln 2}$ | Applies $t = "2"$ and $m(\mathbf{N}) = \frac{-1}{m(\mathbf{T})}$ | | M1 |
| | $y - 3 = \frac{1}{8 \ln 2} (x - 0)$ or $y = 3 + \frac{1}{8 \ln 2} x$ or equivalent. | | | M1 A1 oe cso |
| | | (5) | | |
| 9(d) | $\text{Area}(R) = \int (2^t - 1) \cdot \left(-\frac{1}{2}\right) dt$ | Complete substitution for both y and dx | M1 | |
| | $x = -1 \rightarrow t = 4$ and $x = 1 \rightarrow t = 0$ | | B1 | |
| | $= \left\{ -\frac{1}{2} \right\} \left(\frac{2^t}{\ln 2} - t \right)$ Either $2^t \rightarrow \frac{2^t}{\ln 2}$ or $(2^t - 1) \rightarrow \frac{(2^t)}{\pm \alpha (\ln 2)} - t$ or $(2^t - 1) \rightarrow \pm \alpha (\ln 2)(2^t) - t$ | | | M1* |
| | $(2^t - 1) \rightarrow \frac{2^t}{\ln 2} - t$ | | | A1 |
| | $\left\{ -\frac{1}{2} \left[\frac{2^t}{\ln 2} - t \right]_4^0 \right\} = -\frac{1}{2} \left(\left(\frac{1}{\ln 2} \right) - \left(\frac{16}{\ln 2} - 4 \right) \right)$ | | | |
| | Depends on the previous method mark. Substitutes their changed limits in t and subtracts either way round. | | | dM1* |

A level Mathematics Practice Paper – Integration – Mark scheme

| Question | Scheme | | Marks |
|----------|---------------------------|--|-------------------|
| | $= \frac{15}{2\ln 2} - 2$ | $\frac{15}{2\ln 2} - 2$ or equivalent. | A1 |
| | | | (6) |
| | | | (16 marks) |

A level Mathematics Practice Paper – Integration – Mark scheme

| Question | Scheme | Marks |
|--------------|--|------------------|
| 10(a) | $A = \int_0^3 \sqrt{(3-x)(x+1)} dx, x = 1 + 2\sin$ | |
| | $\frac{dx}{d\theta} = 2\cos$ | B1 |
| | $\left\{ \int \sqrt{(3-x)(x+1)} dx \text{ or } \int \sqrt{(3+2x-x^2)} dx \right\}$ | |
| | $= \int \sqrt{(3-(1+2\sin\theta))(1+2\sin\theta+1)} 2\cos\theta \{d\theta\}$ | M1 |
| | $= \int \sqrt{(2-2\sin\theta)(2+2\sin\theta)} 2\cos\theta \{d\theta\}$ | |
| | $= \int \sqrt{(4-4\sin^2\theta)} 2\cos\theta \{d\theta\}$ | |
| | $= \int \sqrt{(4-4(1-\cos^2\theta))} 2\cos\theta \{d\theta\} \text{ or } \int \sqrt{4\cos^2\theta} 2\cos\theta \{d\theta\}$ | M1 |
| | $= 4 \int \cos^2\theta d\theta, \{k=4\}$ | A1 |
| | $0 = 1 + 2\sin \text{ or } 1 = 2\sin \text{ or } \sin = \frac{1}{2} = \frac{\pi}{6}$ and $3 = 1 + 2\sin \text{ or } 2 = 2\sin \text{ or } \sin = 1 = \frac{\pi}{2}$ | B1 |
| | | (5) |
| 10(b) | $\left\{ k \int \cos^2\theta d\theta \right\} = \left\{ k \int \left(\frac{1+\cos 2\theta}{2} \right) d\theta \right\}$ | M1 |
| | $= \left\{ k \left(\frac{\theta}{2} + \frac{1}{4} \sin 2\theta \right) \right\}$ | M1 |
| | $\left\{ \text{So } 4 \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \cos^2\theta d\theta = \left[2\theta + \sin 2\theta \right]_{\frac{\pi}{6}}^{\frac{\pi}{2}} \right\}$ $= \left(2\left(\frac{\pi}{2}\right) + \sin\left(\frac{2\pi}{2}\right) \right) - \left(2\left(\frac{\pi}{6}\right) + \sin\left(\frac{2\pi}{6}\right) \right)$ | |
| | $\left\{ = \left(\pi \right) - \left(\frac{\pi}{3} + \frac{\sqrt{3}}{2} \right) \right\} = \frac{4\pi}{3} + \frac{\sqrt{3}}{2}$ | A1 cao cso |
| | | (3) |
| | | (8 marks) |

A level Mathematics Practice Paper – Integration – Mark scheme

| Question | Scheme | Marks |
|--|--|-------------------|
| 11(a) | $x = 3\theta \sin \theta, \quad y = \sec^3 \theta, \quad 0 \leq \theta < \frac{\pi}{2}$ | |
| | {When $y = 8,$ } $8 = \sec^3 \theta \Rightarrow \cos^3 \theta = \frac{1}{8} \Rightarrow \cos \theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{3}$ $k \text{ (or } x) = 3\left(\frac{\pi}{3}\right)\sin\left(\frac{\pi}{3}\right)$ | M1 |
| | so $k \text{ (or } x) = \frac{\sqrt{3}\pi}{2}$ | A1 |
| | | (2) |
| 11(b) | $\frac{dx}{d\theta} = 3\sin \theta + 3\theta \cos \theta$ | B1 |
| | $\left\{ \int y \frac{dx}{d\theta} \{d\theta\} \right\} = \int (\sec^3 \theta)(3\sin \theta + 3\theta \cos \theta) \{d\theta\}$ | M1 |
| | $= 3 \int \theta \sec^2 \theta + \tan \theta \sec^2 \theta \, d\theta$ | A1 * |
| | $x = 0 \text{ and } x = k \Rightarrow \underline{\alpha = 0} \text{ and } \underline{\beta = \frac{\pi}{3}}$ | B1 |
| | | (4) |
| 11(c) | $\left\{ \int \theta \sec^2 \theta \, d\theta \right\} = \theta \tan \theta - \int \tan \theta \{d\theta\}$ | M1 |
| | | dM1 |
| | $= \theta \tan \theta - \ln(\sec \theta) \quad \text{or} \quad = \theta \tan \theta + \ln(\cos \theta)$ | A1 |
| | $\left\{ \int \tan \theta \sec^2 \theta \, d\theta \right\} = \frac{1}{2} \tan^2 \theta \text{ or } \frac{1}{2} \sec^2 \theta$ or $\frac{1}{2u^2}$ where $u = \cos \theta$ or $\frac{1}{2}u^2$ where $u = \tan \theta$ | M1 |
| | | A1 |
| | $\{\text{Area}(R)\} = \left[3\theta \tan \theta - 3\ln(\sec \theta) + \frac{3}{2} \tan^2 \theta \right]_0^{\frac{\pi}{3}} \text{ or } \left[3\theta \tan \theta - 3\ln(\sec \theta) + \frac{3}{2} \sec^2 \theta \right]_0^{\frac{\pi}{3}}$ | |
| | $= \left(3\left(\frac{\pi}{3}\right)\sqrt{3} - 3\ln 2 + \frac{3}{2}(3) \right) - (0) \text{ or } \left(3\left(\frac{\pi}{3}\right)\sqrt{3} - 3\ln 2 + \frac{3}{2}(4) \right) - \left(\frac{3}{2}\right)$ | |
| $= \frac{9}{2} + \sqrt{3}\pi - 3\ln 2 \text{ or } \frac{9}{2} + \sqrt{3}\pi + 3\ln\left(\frac{1}{2}\right) \text{ or } \frac{9}{2} + \sqrt{3}\pi - \ln 8 \text{ or}$ $\ln\left(\frac{1}{8}e^{\frac{9}{2} + \sqrt{3}\pi}\right)$ | A1 o.e. | |
| | | (6) |
| | | (12 marks) |

A level Mathematics Practice Paper – Integration – Mark scheme

| Question | Scheme | | Marks |
|---|---|---|--------------------|
| 12(a) | $1 = A(5 - P) + BP$ | Can be implied. | M1 |
| | $A = \frac{1}{5}, B = \frac{1}{5}$ | Either one. | A1 |
| | giving $\frac{\frac{1}{5}}{P} + \frac{\frac{1}{5}}{(5 - P)}$ | | A1 cao, aef |
| | | | (3) |
| 12(b) | $\int \frac{1}{P(5 - P)} dP = \int \frac{1}{15} dt$ | | B1 |
| | $\frac{1}{5} \ln P - \frac{1}{5} \ln(5 - P) = \frac{1}{15}t (+ c)$ | | M1* |
| | $\{t = 0, P = 1 \Rightarrow\} \frac{1}{5} \ln 1 - \frac{1}{5} \ln(4) = 0 + c \quad \left\{ \Rightarrow c = -\frac{1}{5} \ln 4 \right\}$ | | dM1* |
| | e.g.: $\frac{1}{5} \ln \left(\frac{P}{5 - P} \right) = \frac{1}{15}t - \frac{1}{5} \ln 4$ | Using any of the subtraction (or addition) laws for logarithms CORRECTLY | dM1* |
| | $\ln \left(\frac{4P}{5 - P} \right) = \frac{1}{3}t$ | | |
| | e.g.: $\frac{4P}{5 - P} = e^{\frac{1}{3}t}$ or e.g.: $\frac{5 - P}{4P} = e^{-\frac{1}{3}t}$ | Eliminate ln's correctly. | dM1* |
| | gives $4P = 5e^{\frac{1}{3}t} - Pe^{\frac{1}{3}t} \Rightarrow P(4 + e^{\frac{1}{3}t}) = 5e^{\frac{1}{3}t}$ | | |
| | $P = \frac{5e^{\frac{1}{3}t}}{(4 + e^{\frac{1}{3}t})} \quad \left\{ \begin{array}{l} (\div e^{\frac{1}{3}t}) \\ (\div e^{\frac{1}{3}t}) \end{array} \right\}$ | Make P the subject. | dM1* |
| $P = \frac{5}{(1 + 4e^{-\frac{1}{3}t})}$ or $P = \frac{25}{(5 + 20e^{-\frac{1}{3}t})}$ etc. | | A1 | |
| | | | (8) |
| 12(c) | $1 + 4e^{-\frac{1}{3}t} > 1 \Rightarrow P < 5$. So population cannot exceed 5000. | | B1 |
| | | | (1) |
| | | | (12 marks) |

A level Mathematics Practice Paper – Integration – Mark scheme

| | Source paper | Question number | New spec references | Question description | New AOs |
|----|--------------|-----------------|---------------------------|--|----------------------|
| 1 | C4 2012 | 1 | 2.10, 8.6 | Partial fractions, Integration | 1.1b |
| 2 | C4 Jan 2011 | 1 | 8.4, 5.3 | Integration | 1.1b |
| 3 | C4 Jan 2013 | 2 | 8.2, 8.5 | Integration | 1.1b |
| 4 | C4 Jan 2012 | 2 | 8.2, 8.5 | Integration | 1.1b, |
| 5 | C4 2015 | 3 | 6.3, 8.5 | Integration | 1.1b, 2.1, 3.1a |
| 6 | C4 2013 | 3 | 8.2, 8.5, 6.4 | Integration | 1.1b, 2.1 |
| 7 | C4 Jan 2011 | 3 | 2.10, 8.6, 8.7 | Algebra and functions, Integration | 1.1b, 3.1a |
| 8 | C4 2016 | 6 | 8.5, 8.6 | Integration | 1.1b, 2.1, 3.1a |
| 9 | C4 Jan 2013 | 5 | 3.3, 7.3, 7.5, 8.3, 8.5 | Parametric curves and equations, Parametric differentiation, Integration | 1.1b, 3.1a |
| 10 | C4 2015 | 6 | 8.2, 8.3, 8.5 | Integration | 1.1b, 2.1 |
| 11 | C4 2017 | 8 | 5.4, 7.4, 8.3, 8.5 | Parametric curves and equations, Integration | 1.1b, 2.1, 3.1a |
| 12 | C4 Jan 2012 | 8 | 2.10, 8.6, 8.7, 8.8, 2.11 | Partial fractions, Integration | 1.1b, 2.1, 3.1a, 3.4 |